



<b>Test Statistic calculation</b>	<b>Identify the following:</b> <ul style="list-style-type: none"> <li>• Sample means: degrees of freedom</li> <li>• Proportions: show proportions, combined proportion <b>(Ideally, show these values in the formula)</b></li> </ul> <ul style="list-style-type: none"> <li>• z or t test statistic</li> </ul>
<b>Obtain P-value</b>	<b>If using the calculator:</b> normCDF, tCDF, etc. - show values and identify each value.  The two sample t test gives t= 1.27, d.f= 11.7 and p-value of 0.11. This will earn you full credit on the Do part if the numbers are right. There is no chance to earn partial credit when reporting this if any numbers are wrong in the calculator.  Recommended to sketch graph and label findings. Label sketch with mean 0 and the z-score. Show the shaded region with the p-value labeled.
<b>Make a decision</b>	<ul style="list-style-type: none"> <li>• <math>P &lt; \alpha</math> Reject the null hypothesis, statistically significant</li> <li>• <math>P \geq \alpha</math> Fail to reject null hypothesis, not statistically significant</li> </ul> <p>*Using the phrase "accept Ho" gets a point deduction on the AP exam</p>
<b>State conclusion</b>	<ul style="list-style-type: none"> <li>• Assuming Ho is true....</li> <li>• Interpret P: The probability of getting a value from the sample as extreme as the one observed.</li> <li>• Because our p-value is (less than, greater than alpha we do/do not have evidence</li> <li>• Findings are statistically significant at the alpha=</li> <li>• Conclusion in context.</li> </ul>

\*If you attempt to answer a question in two different ways then you get graded on the worse of the two responses.

\*If you show correct calculator work with wrong supplemental work then you will get graded as if the calculator work was not there.

**Commonly used formulas**

Standard Deviation Known  $z^* = \text{invnorm}\left(\frac{1-c}{2}\right)$

Means:  $n \geq z^* \left(\frac{\sigma}{me}\right)^2$

Standard Deviation Unknown  $t^* = \text{invT}\left(\frac{1-c}{2}\right), df$

**P-values:**

Proportions: normcdf (LB, test statistic)

Means: normcdf or tcdf (t, UB, DF)

**Inference Procedure Template-Confidence Intervals**

Panic  
 Parameter  
 Assumptions  
 Name the interval  
 Interval  
 Conclusion in context

<b>Parameter of Interest</b>	Support your reason for choosing the interval you are choosing by mentioning the parameter of interest, the confidence level and the populations.
<b>Conditions of Test or Assumptions</b> <ul style="list-style-type: none"> <li>• Random</li> <li>• Normality</li> <li>• Independence</li> </ul>	Show actual counts and comparisons. <ul style="list-style-type: none"> <li>• Name the condition you are addressing.</li> <li>• <b>Normality:</b> If data is given, you must graph it and check for skewness and outliers; pop. distribution is Normal or <math>n \geq 30</math> (for means)</li> <li>• <b>Normality:</b> Best to show-  <math>n(\hat{p})=150(0.32)=48 \geq 10</math> and <math>n(1-\hat{p})=150(1-.32)=102 \geq 10</math> (for proportions)            If you are not sure about p, <math>p_0</math>, <math>\hat{p}</math>-hat then just show the numbers for partial credit.</li> <li>• <b>Independence:</b> Pop <math>\geq 10n</math></li> </ul>
<b>Name the Interval</b>	<b>Name the interval</b> 2 proportion z interval, 1 proportion z-interval 2 sample t interval, 1 sample t-interval Or <b>Show Formula</b> Only show the formula with the variables if you are absolutely sure that you know the symbols. Don't confuse p with $\hat{p}$ or $p_0$
<b>Test Statistic calculation</b>	Ideally, show formula with values substituted in to it. Then state the Interval endpoints (lower bound-upper bound)  <b>If using the calculator:</b> normCDF, tCDF, ect. -show values and identify each value  The two-proportion z interval gives us the interval - .12234 to .3454. This will earn you full credit on the Do part if the numbers are right. There is no chance to earn partial credit when reporting this if any numbers are wrong in the calculator.
<b>State conclusion</b>	We can be 95% confidence that the interval from ____ to ____ captures the true parameter of the population (in context).